

Monoclonal Mouse Antibody to Human Glycophorin C

Catalog No.:	Mob 067, Mob 067-05
Intended Use:	This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy. Clinical interpretation of staining results should be accompanied by histological studies with proper controls. Patients' clinical histories and other relevant diagnostic tests should be utilized by a qualified person(s) when evaluating and interpreting results.
Immunogen:	BALB/C mice were injected with Red cell ghosts.
Clone:	Ret40f
Isotype:	IgG1, Kappa
Format:	This antibody is supplied as tissue culture supernatant containing sodium azide as a preservative.
Titer/Working Dilution:	This antibody may be diluted to a titer of 1:15-1:30 in an ABC method. The final dilution should be determined by the user based upon the staining conditions employed.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed. <u>Suitable for formalin fixed, paraffin embedded tissue sections.</u>
Specificity:	This antibody is specific to beta sialoglycophorin also known as Glycophorin C. It is found on the human erythrocyte membrane and is one of four sialic acid rich polypeptide.
Positive Control:	Tonsil
Cellular Localization:	Cell membrane
Storage:	Store at 2-8°C. Do not use beyond the expiration date stated on the label.
References:	i) Gatter et al. Histopathol 13: 257, 1988.

IVD: For In Vitro Diagnostic Use

DBS will not be held responsible for patent infringement or other violation that may occur with the use of our product

DBS

1020 Serpentine Lane, # 114, Pleasanton, CA 94566 Tel: 925 484 3350, Fax: 925 484 3390

Website: www.dbiosys.com e-mail: customersupport@dbiosys.com