

Monoclonal Mouse Antibody to Human Glycophorin A

Catalog No.:	PDM 124
Intended Use:	This product is intended for qualitative immunohistochemistry with normal and neoplastic formalin-fixed, paraffin-embedded tissue sections, to be viewed by light microscopy. Clinical interpretation of staining results should be accompanied by histological studies with proper controls. Patients' clinical histories and other relevant diagnostic tests should be utilized by a qualified person(s) when evaluating and interpreting results.
Immunogen:	BALB/C mice were immunized with formalin fixed membranefrom a Hairy cell leukemia.
Clone:	JC159
Isotype:	IgG1, Kappa
Format:	This antibody has been pretitered and quality controlled to work on formalin-fixed paraffin-embedded and acetone fixed cryostat tissue sections. No further titration is required.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed. <u>Suitable for formalin fixed, paraffin embedded tissue sections.</u>
Positive Control:	Tonsil
Cellular Localization:	Cell membrane
Specificity:	This antibody is specific to erythroid cells at various stages of differentiation from the erythroblast to the mature red cell.
Storage:	Store at 2-8°C. Do not use beyond the expiration date stated on the label.
References:	i) Erber et al. Hematol Rev %: 113, 1991. ii) Greaves et al. Blood 61: 645, 1983. iii) Loken et al. Blood 69: 255, 1987.

IVD: For In Vitro Diagnostic Use

DBS will not be held responsible for patent infringement or other violation that may occur with the use of our product

DBS

1020 Serpentine Lane, # 114, Pleasanton, CA 94566 Tel: 925 484 3350, Fax: 925 484 3390

Website: www.dbiosys.com e-mail: customersupport@dbiosys.com