

Rabbit Polyclonal Antibody to Apoptosis Protease Activating Factor 1 (Apaf-1)

Catalog No.:	RP 136, RP 136-05
Applications:	Research Use Only. Optimal working conditions must be determined by the end user.
Immunogen:	A synthetic peptide derived from C-terminus of human Apaf-1.
Host:	Rabbit
Format:	Purified immunoglobulin fraction of rabbit antiserum against Apaf-1 containing sodium azide as a preservative.
Titer/Working Dilution:	This antibody may be diluted to a titer of 1:50-1:100 in an ABC method. The final dilution should be determined by the user based upon the staining conditions employed.
Staining Protocol:	We suggest an incubation period of 30 minutes at room temperature. Optimal incubation conditions should be determined by the user based upon the fixation conditions and staining system employed. <u>Formalin fixed paraffin embedded tissue sections require high temperature antigen unmasking with 10 mM citrate buffer, pH 6.0 prior to immunostaining.</u>
Specificity:	Apaf-1 is a cytosolic protein that rests in a latent state until bound by cytochrome c (Apaf-2). The resulting Apaf-1/cytochrome c complex associates with the zymogen form of caspase-9 (Apaf-3) in the presence of dATP or ATP, promoting the autocatalytic activation of caspase-9. This antibody is useful in studies of the activation of caspases through Apaf-1 and also in metastatic melanomas where Apaf-1 is inactivated which leads to defects in the execution of apoptotic cell death. This antibody cross reacts with mouse and rat.
Positive Control:	Testis
Cellular Localization:	Cytoplasmic
Storage:	Store at 2-8°C. Do not use beyond the expiration date stated on the label.

RUO: Research Use Only

DBS will not be held responsible for patent infringement or other violation that may occur with the use of our product

DBS

1020 Serpentine Lane, # 114, Pleasanton, CA 94566 Tel: 925 484 3350, Fax: 925 484 3390

Website: www.dbiosys.com e-mail: customersupport@dbiosys.com